



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

From these equations we find the values of A, B, C, D, E ; whence,

$$U_n = -20 + 36n - \frac{115n^2}{6} + \frac{9n^3}{2} - \frac{n^4}{3}.$$

From this we get the 6th term to be 46.

Thus, the series is

$$1 + 6 + 10 + 20 + 35 + 46 + \dots$$

Using the method of differences we get

$$\begin{array}{ccccccc} 5 & 4 & 10 & 15 & 11 & \dots \\ -1 & 6 & 5 & -4 & \dots & \\ 7 & -1 & -9 & \dots & \dots & \\ -8 & -8 & \dots & \dots & \dots & \\ 0 & \dots & \dots & \dots & \dots & \end{array}$$

Then

$$\begin{aligned} S_n &= n + \frac{5n(n-1)}{|2|} - \frac{n(n-1)(n-2)}{|3|} + \frac{7n(n-1)(n-2)(n-3)}{|4|} - \frac{8n(n-1)(n-2)(n-3)(n-4)}{|5|} \\ &= -\frac{n}{120} (8n^4 - 115n^3 + 510n^2 - 1145n + 622). \end{aligned}$$

(c) In the series $1 + 5 + 15 + 35 + 70 + \dots$

$$U_n = A + Bn + Cn^2 + Dn^3 + En^4 + \dots$$

$$\left\{ \begin{array}{l} A + B + C + D + E = 1, \\ A + 2B + 4C + 8D + 16E = 5, \\ A + 3B + 9C + 27D + 81E = 15, \\ A + 4B + 16C + 64D + 256E = 35, \\ A + 5B + 25C + 125D + 625E = 70, \end{array} \right\} \begin{array}{l} A = 0, \\ B = 1/4, \\ C = 11/24, \\ D = 1/4, \\ E = 1/24. \end{array}$$

Hence,

$$U_n = \frac{n}{4} + \frac{11n^2}{24} + \frac{n^3}{4} + \frac{n^4}{24}.$$

Then

$$24S_n = 6\Sigma n + 11\Sigma n^2 + 6\Sigma n^3 + \Sigma n^4,$$

$$24S_n = 3n(n+1) + \frac{11n(n+1)(2n+1)}{6} + \frac{3n^2(n+1)^2}{2} + \frac{n(n+1)(6n^3+9n^2+n-1)}{30}.$$

$$S_n = \frac{n}{120} (n^4 + 10n^3 + 35n^2 + 50n + 24) = \frac{n}{120} (n+1)(n+2)(n+3)(n+4).$$

Also solved by O. S. ADAMS, H. H. CONWELL, PAUL CAPRON, and WILLIAM TIER.

GEOMETRY.

496. Proposed by NATHAN ALTSHILLER, University of Oklahoma.

Find all the lines such that the pairs of tangent planes to a given sphere (ellipsoid) passing through them, shall be orthogonal.

SOLUTION BY WILLIAM HOOVER, Columbus, Ohio.

One obvious solution is given when one of the tangent planes is fixed in position, for then this plane is the locus of lines common to it and a second tangent plane orthogonal to the first.

It is not troublesome to show that the equation of a first plane embracing a line

$$\frac{x-a}{l} = \frac{y-b}{m} = \frac{z-c}{n}, \quad (1)$$

$$\frac{(x-a)\lambda_1}{l} + \frac{(y-b)\mu_1}{m} + \frac{(z-c)\nu_1}{n} = 0, \quad (2)$$

with the condition

$$\lambda_1 + \mu_1 + \nu_1 = 0, \quad (3)$$

and that of a second plane embracing (1) is

$$\frac{(x-a)\lambda_2}{l} + \frac{(y-b)\mu_2}{m} + \frac{(z-c)\nu_2}{n} = 0, \quad (4)$$

with the condition

$$\lambda_2 + \mu_2 + \nu_2 = 0. \quad (5)$$

If (2) and (4) are orthogonal,

$$\frac{\lambda_1}{l} \cdot \frac{\lambda_2}{l} + \frac{\mu_1}{m} \cdot \frac{\mu_2}{m} + \frac{\nu_1}{n} \cdot \frac{\nu_2}{n} = 0, \quad (6)$$

and if (2) and (4) touch

$$a_1x^2 + b_1y^2 + c_1z^2 = 1, \quad (7)$$

$$\left(\frac{\lambda_1}{l}\right)^2/a_1 + \left(\frac{\mu_1}{m}\right)^2/b_1 + \left(\frac{\nu_1}{n}\right)^2/c_1 = \left(\frac{a\lambda_1}{l} + \frac{b\mu_1}{m} + \frac{c\nu_1}{n}\right)^2, \quad (8)$$

and

$$\left(\frac{\lambda_2}{l}\right)^2/a_1 + \left(\frac{\mu_2}{m}\right)^2/b_1 + \left(\frac{\nu_2}{n}\right)^2/c_1 = \left(\frac{a\lambda_2}{l} + \frac{b\mu_2}{m} + \frac{c\nu_2}{n}\right)^2. \quad (9)$$

Now (3), (5), (6), (8), (9) is a system of *five* equations for the determination of the *six* ratios $\lambda_1/\nu_1, \mu_1/\nu_1; \lambda_2/\nu_2, \mu_2/\nu_2$; and $l/n, m/n$, giving an indeterminate solution. The values of $l/n, m/n$ are the only ones needed in (1).

There seems to be a missing condition in the statement of the problem.

If the direction of the line (1) were constant, or $l : m : n$, constant, the *locus* of the line (1) would be a right circular cylinder, or, in other words, the locus of the line of intersection of constant direction of pairs of orthogonal tangent planes to a central conicoid is a right circular cylinder.

Note.—The single condition on the line restricts it to be a member of a line complex whose order apparently may be as high as 8. It would be desirable to determine this explicitly. EDITORS.

497. Proposed by NATHAN ALTSHILLER, University of Oklahoma.

Find the locus of the mid-point of the segment determined by two spheres on any line passing through any point common to the two spheres.

SOLUTION BY S. W. REAVES, University of Oklahoma.

Let the plane of the common circle of the spheres be chosen for yz -plane, and the line of centers for x -axis. Let the radius of the common circle be k , and let $(a, 0, 0)$ and $(b, 0, 0)$ be the centers of the two spheres. Then the equation of one sphere is

$$(x-a)^2 + y^2 + z^2 = a^2 + k^2, \quad (1)$$

or

$$x^2 + y^2 + z^2 - 2ax = k^2;$$

and, similarly, the equation of the other is

$$x^2 + y^2 + z^2 - 2bx = k^2. \quad (2)$$

Let l, m, n be the direction cosines of an arbitrary line through the point $M(0, 0, k)$. Then the equation of the line may be written

$$\frac{x}{l} = \frac{y}{m} = \frac{z-k}{n} = r, \quad (3)$$

where r is the length of the segment joining $(0, 0, k)$ and (x, y, z) .

To find the length of the segment MP cut from this line by the sphere (1), we substitute $x = lr, y = mr, z = nr + k$ in equation (1) and solve for that value of r which is not zero. We thus find at once

$$r = MP = 2al - 2kn.$$

Substituting the same values in (2), we find likewise for the segment MQ intercepted by the